

# BOUNDARY INTEGRAL OPERATOR FOR THE FRACTIONAL LAPLACIAN IN THE BOUNDED SMOOTH DOMAIN

TONGKEUN CHANG

**ABSTRACT.** We study the boundary integral operator induced from the fractional Laplace equation in a bounded smooth domain. For  $\frac{1}{2} < \alpha < 1$ , we show the bijectivity of the boundary integral operator  $S_{2\alpha} : L^p(\partial\Omega) \rightarrow H_p^{2\alpha-1}(\partial\Omega)$ ,  $1 < p < \infty$ . As an application, we show the existence of the solution of the boundary value problem of the fractional Laplace equation.

## 1. INTRODUCTION

In this paper, we study a boundary integral operator defined on the boundary of a bounded domain in  $\mathbb{R}^n$ ,  $n \geq 3$ . Let  $\Omega$  be a bounded smooth domain and  $\Gamma_{2\alpha}(x) = c(n, 2\alpha) \frac{1}{|x|^{n-2\alpha}}$  be the Riesz kernel of order  $2\alpha$ ,  $0 < 2\alpha < n$  in  $\mathbb{R}^n$ , where  $c(n, 2\alpha)$  is the normalized constant. The layer potential of a fractional Laplacian for  $\phi \in L^p(\partial\Omega)$ ,  $1 < p < \infty$  is defined by

$$\mathcal{S}_{2\alpha}\phi(x) = \int_{\partial\Omega} \Gamma_{2\alpha}(x - Q)\phi(Q)dQ, \quad x \in \mathbb{R}^n \setminus \partial\Omega. \quad (1.1)$$

The boundary integral operator

$$S_{2\alpha}\phi = (\mathcal{S}_{2\alpha}\phi)|_{\partial\Omega} \quad (1.2)$$

is defined by the restriction of  $\mathcal{S}_{2\alpha}\phi$  over  $\partial\Omega$ .

M. Zähle studied the Riesz potentials in a general metric space  $(X, \rho)$  with Ahlfors  $d$ -regular measure  $\mu$ . She showed  $S_{2\alpha} : L^2(X, d\mu) \rightarrow L_{2\alpha}^2(X, d\mu)$ ,  $0 < 2\alpha < n$  is invertible, where  $L^2(X, d\mu)$  is decomposed by the null space  $N(S_{2\alpha})$  and the orthogonal complement of  $N(S_{2\alpha})$ , that is,  $L^2(X, d\mu) = N(S_{2\alpha}) \otimes L_{2\alpha}^2(X, d\mu)$  (see [19] and [20]).

The author of [5] showed that the boundary integral operator  $S_{2\alpha}$  defined in (1.2) is extended to  $H_2^{-\alpha+\frac{1}{2}}(\partial\Omega)$ ,  $\frac{1}{2} < \alpha < 1$  such that  $S_{2\alpha} : H_2^{-\alpha+\frac{1}{2}}(\partial\Omega) \rightarrow H_2^{\alpha-\frac{1}{2}}(\partial\Omega)$  is bijective operator and  $\mathcal{S}_{2\alpha}\phi \in \dot{H}_2^\alpha(\mathbb{R}^n)$  for  $\phi \in H_2^{-\alpha+\frac{1}{2}}(\partial\Omega)$  (see section 2 for the definition of the function spaces).

When  $2\alpha = 2$ ,  $\Gamma_2$  is the fundamental solution of the Laplace equation in  $\mathbb{R}^n$  and (1.1) is a single layer potential of the Laplace equation. The single layer potential and boundary layer potential of the Laplace equation were studied by many mathematicians to show the existence of the solution of the boundary value problem of the Laplace equation in a bounded domain (see [9], [11], [13] and [18]).

---

1991 *Mathematics Subject Classification.* Primary 45P05 ; Secondary 30E25.

*Key words and phrases.* boundary integral operator, layer potential, fractional Laplacian, bounded domain.

The first result of this paper is the following Theorem.

**Theorem 1.1.** *Let  $\Omega$  be a bounded  $C^2$ -domain in  $\mathbb{R}^n$ ,  $3 \leq n$ . Let  $\frac{1}{2} < \alpha < 1$  and  $1 < p < \infty$ . Then,  $S_{2\alpha} : L^p(\partial\Omega) \rightarrow H_p^{2\alpha-1}(\partial\Omega)$  is bijective.*

The function space  $H_p^{2\alpha-1}(\partial\Omega)$  is defined in the section 2.

The layer potential for  $\phi \in B_p^s(\partial\Omega)$ ,  $s < 0$ ,  $1 < p < \infty$  is defined by

$$\mathcal{S}_{2\alpha}\phi(x) = \langle \phi, \Gamma_{2\alpha}(x - \cdot) \rangle, \quad x \in \mathbb{R}^n \setminus \partial\Omega, \quad (1.3)$$

where  $\langle \cdot, \cdot \rangle$  is the duality pairing between  $B_p^s(\partial\Omega)$  and  $B_{p'}^{-s}(\partial\Omega)$ ,  $\frac{1}{p} + \frac{1}{p'} = 1$ . In particular, if  $\phi \in L^p(\partial\Omega)$ , then  $\mathcal{S}_{2\alpha}\phi$  is defined by (1.1). The second result is the following Theorem.

**Theorem 1.2.** *Let  $\frac{1}{2} < \alpha < 1$  and  $1 < p < \infty$ . For  $\phi \in B_p^s(\partial\Omega)$ , let  $u = \mathcal{S}_{2\alpha}\phi$  be a layer potential defined in (1.3). Let  $-2\alpha + 1 - \frac{1}{p} < s < 0$ . Then,  $u \in B_{loc,p}^{s+2\alpha-1+\frac{1}{p}}(\mathbb{R}^n)$  and*

$$\|u\|_{B_p^{s+2\alpha-1+\frac{1}{p}}(B_R)} \leq c_R \|\phi\|_{B_p^s(\partial\Omega)}, \quad (1.4)$$

where  $B_R$  is open ball in  $\mathbb{R}^n$  whose radius is  $R$  and center is origin such that  $\Omega \subset B_R$ . Moreover, if  $p > \frac{n-1}{n+s-1}$ , the  $u \in \dot{B}_p^{s+2\alpha-1+\frac{1}{p}}(\mathbb{R}^n)$  such that

$$\|u\|_{\dot{B}_p^{s+2\alpha-1+\frac{1}{p}}(\mathbb{R}^n)} \leq c \|\phi\|_{B_p^s(\partial\Omega)}. \quad (1.5)$$

The function spaces  $B_{loc,p}^{s+2\alpha-1+\frac{1}{p}}(\mathbb{R}^n)$  and  $\dot{B}_p^{s+2\alpha-1+\frac{1}{p}}(\mathbb{R}^n)$  are defined in the section 2.

The boundary integral operators (the single layer potential and the double layer potential) have been studied by many mathematicians. The bijectivity of the layer potential has been used to show the existence of the solutions of the partial differential equations in a bounded domain or bounded cylinder (see [3], [4], [7], [10], [12], [15] and [16]).

As in the vast literature, we apply the bijectivity of the boundary integral operator to the boundary value problem of the fractional Laplace equation in the bounded smooth domain. The fractional Laplacian of order  $0 < \alpha < 1$  of a function  $v : \mathbb{R}^n \rightarrow \mathbb{R}$  is expressed by the formula

$$\Delta^\alpha v(x) = C(n, \alpha) \int_{\mathbb{R}^n} \frac{v(x+y) - 2v(x) + v(x-y)}{|x-y|^{n+2\alpha}} dy,$$

where  $C(n, \alpha)$  is some normalization constant. The fractional Laplacian can also be defined as a pseudo-differential operator

$$\widehat{(-\Delta)^\alpha v}(\xi) = (2\pi|\xi|)^{2\alpha} \hat{v}(\xi),$$

where  $\hat{v}(\xi) := \int_{\mathbb{R}^n} v(x) e^{-2\pi i \xi \cdot x} dx$ ,  $\xi \in \mathbb{R}^n$  is the Fourier transform of  $v$  in  $\mathbb{R}^n$ . In particular, when  $2\alpha = 2$  it is the Laplacian  $\Delta v(x) = \sum_{1 \leq i \leq n} \frac{\partial^2}{\partial x_i^2} v(x)$ .

**Definition 1.3.** Let  $0 < \alpha < 1$ . We say that  $v$  is a weak solution of  $\Delta^\alpha u = 0$  in  $\mathbb{R}^n \setminus \partial\Omega$  if  $v$  satisfies

$$\int_{\mathbb{R}^n} v(x) \Delta^\alpha \psi(x) dx = \int_{\mathbb{R}^n} (2\pi|\xi|)^{2\alpha} \hat{v}(\xi) \bar{\hat{\psi}}(\xi) d\xi = 0 \quad (1.6)$$

for all  $\psi \in C_c^\infty(\mathbb{R}^n \setminus \partial\Omega)$ .

In fact, if  $u$  is weak solution, then  $u$  is continuous function in  $\mathbb{R}^n \setminus \partial\Omega$  and satisfies

$$\Delta^\alpha u(x) = 0 \quad \text{for} \quad x \in \mathbb{R}^n \setminus \partial\Omega.$$

(See Theorem 3.9 in [2]).

For the application of Theorem 1.1 and Theorem 1.2, we show the existence of the solution of the boundary value problem of the fractional Laplace equation in the bounded smooth domain.

**Theorem 1.4.** Let  $\Omega$  be a bounded  $C^2$  domain in  $\mathbb{R}^n$ ,  $3 \leq n$  and  $\frac{1}{2} < \alpha < 1$ . Let  $0 < t < 1$  and  $1 < p < \infty$ . Then, for given  $g \in B_p^t(\partial\Omega)$ , the following equation

$$\begin{cases} \Delta^\alpha u = 0 & \text{in } \mathbb{R}^n \setminus \partial\Omega, \\ u|_{\partial\Omega} = g \in B_p^t(\partial\Omega), \\ u \in B_{loc,p}^{t+\frac{1}{p}}(\mathbb{R}^n), \\ |u(x)| = O(|x|^{-n+2\alpha}) \quad \text{near } \infty. \end{cases} \quad (1.7)$$

has a weak solution. And, if  $\frac{n-1}{n+t-2\alpha} < p < \infty$ , then  $u \in \dot{B}_p^{t+\frac{1}{p}}(\mathbb{R}^n)$ . Furthermore, there exists  $\phi \in B_p^{t-2\alpha+1}(\partial\Omega)$  such that

$$u = \mathcal{S}_{2\alpha} \phi. \quad (1.8)$$

The rest of this paper is organized as follows. In section 2, we introduce the several function spaces. In section 3, we introduce the several property of the layer potential. In section 4, we prove the Theorem 1.1. In section 5, we prove the Theorem 1.2. In section 6, we prove the Theorem 1.4.

## 2. FUNCTION SPACES

**2.1. Function spaces in  $\mathbb{R}^n$ .** In this section, we introduce the several function spaces, Sobolev space and Besov space. For  $s \in \mathbb{R}$ , we consider a distribution  $G_s$  whose Fourier transform in  $\mathbb{R}^n$  is defined by

$$\widehat{G_s}(\xi) = (1 + 4\pi^2|\xi|^2)^{-\frac{s}{2}}.$$

For  $s \in \mathbb{R}$ ,  $1 \leq p \leq \infty$ , we define the Sobolev space  $H_p^s(\mathbb{R}^n)$  by

$$H_p^s(\mathbb{R}^n) : = \{f \in \mathcal{S}'(\mathbb{R}^n) \mid \|f\|_{H_p^s(\mathbb{R}^n)} := \|G_{-s} * f\|_{L^p(\mathbb{R}^n)} < \infty\},$$

where  $*$  is a convolution in  $\mathbb{R}^n$  and  $\mathcal{S}'(\mathbb{R}^n)$  is the dual space of the Schwartz space  $\mathcal{S}(\mathbb{R}^n)$ . In particular, when  $s = k \in \mathbb{N} \cup \{0\}$  and  $1 < p < \infty$ ,

$$H_p^k(\mathbb{R}^n) = \{f \mid D^\beta f \in L^p(\mathbb{R}^n), \quad |\beta| \leq k\},$$

where  $\beta \in (\mathbb{N} \cup \{0\})^n$  and  $|\beta| = \beta_1 + \beta_2 + \cdots + \beta_n$  for  $\beta = (\beta_1, \beta_2, \dots, \beta_n)$ .

For  $k < s < k+1$ ,  $k \in \mathbb{N}$ , we define the Besov space  $B_p^s(\mathbb{R}^n)$  and homogeneous Besov space  $\dot{B}_p^s(\mathbb{R}^n)$  by

$$B_p^s(\mathbb{R}^n) = \{f \in \mathcal{S}'(\mathbb{R}^n) \mid \|f\|_{B_p^s} < \infty\}, \quad \dot{B}_p^s(\mathbb{R}^n) = \{f \in \mathcal{S}'(\mathbb{R}^n) \mid \|f\|_{\dot{B}_p^s} < \infty\}$$

with the norms

$$\begin{aligned} \|f\|_{B_p^s} &:= \|f\|_{H_p^k} + \left( \sum_{|\beta|=k} \int \int_{\mathbb{R}^n \times \mathbb{R}^n} \frac{|D^\beta f(x) - D^\beta f(y)|^p}{|x-y|^{n+p(s-k)}} dy dx \right)^{\frac{1}{p}}, \\ \|f\|_{\dot{B}_p^s} &:= \left( \sum_{|\beta|=k} \int \int_{\mathbb{R}^n \times \mathbb{R}^n} \frac{|D^\beta f(x) - D^\beta f(y)|^p}{|x-y|^{n+p(s-k)}} dy dx \right)^{\frac{1}{p}}. \end{aligned}$$

If  $s \in \mathbb{R}$  is negative, then we define  $B_p^s$  and  $\dot{B}_p^s$  as the dual spaces of  $B_{p'}^{-s}$  and  $\dot{B}_{p'}^{-s}$ , respectively, where  $\frac{1}{p} + \frac{1}{p'} = 1$ . The real interpolation method and complex interpolation method give

$$(H_p^{s_0}, H_p^{s_1})_{\theta, p} = B_p^s, \quad [H_p^{s_0}, H_p^{s_1}]_\theta = H_p^s$$

for  $s = (1-\theta)s_0 + \theta s_1$ ,  $s_0, s_1 \in \mathbb{R}$ ,  $0 < \theta < 1$  (see theorem 6.4.5 in [1]).

**2.2. Function spaces in  $\Omega$ .** Let  $\Omega$  be bounded  $C^2$ -domain in  $\mathbb{R}^n$ . Let  $R_\Omega f$  be a restriction over  $\Omega$  of the function  $f$  defined in  $\mathbb{R}^n$ . For  $s \geq 0$ , we define the function spaces

$$H_p^s(\Omega) := \{R_\Omega f \mid f \in H_p^s(\mathbb{R}^n)\}, \quad B_p^s(\Omega) := \{R_\Omega f \mid f \in B_p^s(\mathbb{R}^n)\}$$

with norms

$$\|f\|_{H_p^s(\Omega)} := \inf \|F\|_{H_p^s(\mathbb{R}^n)}, \quad \|f\|_{B_p^s(\Omega)} := \inf \|F\|_{B_p^s(\mathbb{R}^n)},$$

where infimums are taken in  $F \in H_p^s(\mathbb{R}^n)$  and  $F \in B_p^s(\mathbb{R}^n)$ , respectively, such that  $R_\Omega F = f$ .

Note that for non-negative integer  $k$  and  $1 < p < \infty$ ,

$$H_p^k(\Omega) = \{f \in L^p(\Omega) \mid \sum_{0 \leq l \leq k} \|D^l f\|_{L^p(\Omega)} < \infty\} \quad (2.1)$$

and for  $0 < \theta < 1$ ,

$$(H_p^{k_1}(\Omega), H_p^{k_2}(\Omega))_{\theta, p} = B_p^s(\Omega), \quad [H_p^{k_1}(\Omega), H_p^{k_2}(\Omega)]_\theta = H_p^s(\Omega),$$

where  $s = \theta k_1 + (1-\theta)k_2$  (see chapter 2 in [13]). In particular, for  $k < s < k+1$ , we have

$$\|f\|_{B_p^s(\Omega)} \approx \|f\|_{H_p^k(\Omega)} + \left( \sum_{|\beta|=k} \int_\Omega \int_\Omega \frac{|D_x^\beta f(x) - D_y^\beta f(y)|^p}{|x-y|^{n+p(s-k)}} dx dy \right)^{\frac{1}{p}}.$$

For negative  $s \in \mathbb{R}$ , we define  $B_p^s(\Omega)$  and  $H_p^s(\Omega)$  as the dual spaces of  $B_{p'}^{-s}(\Omega)$  and  $H_{p'}^{-s}(\Omega)$ , respectively, where  $\frac{1}{p} + \frac{1}{p'} = 1$ .

For  $s > 0$ , we define the spaces  $H_{p0}^s(\Omega)$  and  $B_{p0}^s(\Omega)$  as the closures of  $C_c^\infty(\Omega)$  in  $H_p^s(\Omega)$  and  $B_p^s(\Omega)$ , respectively.

**2.3. Function spaces in  $\partial\Omega$ .** Let  $\Omega$  be a bounded  $C^2$ -domain in  $\mathbb{R}^n$ . Let us  $\Delta(P, r) = B(P, r) \cap \partial\Omega$  for  $P \in \partial\Omega$ . Then, there is  $r_0 > 0$  such that for all  $P \in \partial\Omega$ , there is bijective  $C^2$ -function  $\Psi : B'(0, r_0) \rightarrow \Delta(P, r_0)$ , where  $B'(0, r_0)$  is open balls in  $\mathbb{R}^{n-1}$  whose radius is  $R$  and center is origin. Since  $\Omega$  is bounded domain, there are  $P_1, P_2, \dots, P_N$  such that  $\partial\Omega \subset \cup_i \Delta(P_i, r_0)$ . Moreover, there are bijective  $C^2$ -functions  $\Psi_i : B'(0, r_0) \rightarrow \Delta(P_i, r_0)$ . Now, we say that  $\phi$  is in function space  $H_p^s(\partial\Omega)$ ,  $-2 \leq s \leq 2$  if  $\phi \circ \Psi_i \in H_p^s(B'(0, r_0))$  for all  $1 \leq i \leq N$ . And the norm is

$$\|\phi\|_{H_p^s(\partial\Omega)} = \sum_{1 \leq i \leq N} \|\phi \circ \Psi_i\|_{H_p^s(B'(0, r_0))}.$$

Similarly, we define the function space  $B_p^s(\partial\Omega)$ . Clearly, for  $0 < s < 2$ ,  $H_p^{-s}(\partial\Omega)$  and  $B_p^{-s}(\partial\Omega)$  are dual spaces of  $H_{p'}^s(\partial\Omega)$  and  $B_{p'}^s(\partial\Omega)$ ,  $\frac{1}{p} + \frac{1}{p'} = 1$ , respectively.

For  $0 < \theta < 1$ ,

$$(H_p^{k_1}(\partial\Omega), H_p^{k_2}(\partial\Omega))_{\theta, p} = B_p^s(\partial\Omega), \quad [H_p^{k_1}(\partial\Omega), H_p^{k_2}(\partial\Omega)]_\theta = H_p^s(\partial\Omega), \quad (2.2)$$

where  $s = (1 - \theta)k_1 + \theta k_2$  (see chapter 2 in [13]).

### 3. BOUNDARY LAYER POTENTIAL

Let  $S_{2\alpha}$  be a boundary layer potential defined by (1.3). Because of  $\partial\Omega$  is  $C^2$ , we get the following Proposition:

**Proposition 3.1.** *Let  $\Omega$  be  $C^2$  bounded domain. For  $-2 \leq s \leq 3 - 2\alpha$  and  $1 < p < \infty$ ,*

$$S_{2\alpha} : H_p^s(\partial\Omega) \rightarrow H_p^{s+2\alpha-1}(\partial\Omega) \quad (3.1)$$

*is bounded operator. Moreover, for  $-1 \leq s \leq 0$ ,*

$$S_2 : H_p^s(\partial\Omega) \rightarrow H_p^{1+s}(\partial\Omega)$$

*is bijective.*

(See [18]).

Let  $\psi, \phi \in C^2(\partial\Omega)$  and  $S_{2\alpha}^* : H_{p'}^{-s-2\alpha+1}(\partial\Omega) \rightarrow H_{p'}^{-s}(\partial\Omega)$  is a dual operator of (3.1), where  $\frac{1}{p} + \frac{1}{p'} = 1$ . Using (1.1), we have

$$\begin{aligned} \langle\langle S_{2\alpha}^* \psi, \phi \rangle\rangle &= \langle \psi, S_{2\alpha} \phi \rangle = \int_{\partial\Omega} \psi(P) S_{2\alpha} \phi(P) dP \\ &= \int_{\partial\Omega} \phi(P) S_{2\alpha} \psi(P) dP = \langle \phi, S_{2\alpha} \psi \rangle, \end{aligned} \quad (3.2)$$

where  $\langle \cdot, \cdot \rangle$  is the duality pairing between  $H_p^{s+2\alpha-1}(\partial\Omega)$  and  $H_{p'}^{-s-2\alpha+1}(\partial\Omega)$ ,  $\frac{1}{p} + \frac{1}{p'} = 1$ , and  $\langle\langle \cdot, \cdot \rangle\rangle$  is the duality pairing between  $H_p^s(\partial\Omega)$  and  $H_{p'}^{-s}(\partial\Omega)$ ,  $\frac{1}{p} + \frac{1}{p'} = 1$ . Since  $C^2(\partial\Omega)$  is dense subset of  $H_p^s(\partial\Omega)$ , (3.2) implies that  $S_{2\alpha}^* : H_{p'}^{-s-2\alpha+1}(\partial\Omega) \rightarrow H_{p'}^{-s}(\partial\Omega)$  is a same operator with  $S_{2\alpha} : H_{p'}^{-s-2\alpha+1}(\partial\Omega) \rightarrow H_{p'}^{-s}(\partial\Omega)$  for  $s < 0$ .

#### 4. PROOF OF THEOREM 1.1

To prove the Theorem 1.1, we need the following Lemma.

**Lemma 4.1.** *Let  $\frac{1}{2} < \alpha < 1$ . Given sufficiently small  $\epsilon > 0$ , there are bounded linear operators  $T^1 : L^p(\partial\Omega) \rightarrow H_p^1(\partial\Omega)$  with  $\|T^1\|_{L^p(\partial\Omega) \rightarrow H_p^1(\partial\Omega)} < c\epsilon$  and  $T^2 : H_p^{-1}(\partial\Omega) \rightarrow H_p^1(\partial\Omega)$  such that*

$$S_{2\alpha} S_{3-2\alpha} = S_2 + T^1 + T^2. \quad (4.1)$$

**Remark 4.2.** (1) *Since  $S_2 : L^p(\partial\Omega) \rightarrow H_p^1(\partial\Omega)$  is bijective, for sufficiently small  $\epsilon > 0$ ,  $S_2 + T^1 : L^p(\partial\Omega) \rightarrow H_p^1(\partial\Omega)$  is bijective.*

(2) *Since  $S_2, S_{2\alpha} S_{3-2\alpha}$  and  $T^2$  are bounded operators from  $H_p^{-1}(\partial\Omega)$  to  $L^p(\partial\Omega)$ ,  $T^1 : H_p^{-1}(\partial\Omega) \rightarrow L^p(\partial\Omega)$  is also bounded operator. Then, by the complex interpolation property (2.2), we get that for  $-1 < s < 0$ ,*

$$\|T^1\|_{H_p^s(\partial\Omega) \rightarrow H_p^{1+s}(\partial\Omega)} \leq c\epsilon^{1+s}. \quad (4.2)$$

(3) *By the argument of (1), (2) and by Proposition 3.1, we get  $S_2 + T^1 : H_p^s(\partial\Omega) \rightarrow H_p^{1+s}(\partial\Omega)$ ,  $-1 < s < 0$  is bijective.*

**Proof of Lemma 4.1.** Let  $0 < 9\epsilon < r_0$ , where  $r_0 > 0$  is defined in a section 2.3. Let  $P_1, P_2, \dots, P_m \in \partial\Omega$  such that  $|P_i - P_j| > \epsilon$  and  $\partial\Omega \subset \cup_{1 \leq i \leq m} B(P_i, \frac{3}{2}\epsilon)$ . Let  $\{\eta_i\}_{1 \leq i \leq m}$  be a partition of unity of  $\{B(P_i, 2\epsilon)\}_{1 \leq i \leq m}$  such that  $\text{supp } \eta_i \subset B(P_i, 2\epsilon)$ ,  $\eta_i \equiv 1$  in  $B(P_i, \frac{3}{2}\epsilon)$ . Let  $\kappa_i = \sum_{|P_i - P_j| \leq 5\epsilon} \eta_j$  and  $\lambda_i = \sum_{|P_i - P_j| \leq 7\epsilon} \eta_j$  such that  $\text{supp } \kappa_i \subset B(P_i, 7\epsilon)$  and  $\text{supp } \lambda_i \subset B(P_i, 9\epsilon)$ . Then, for  $\phi \in L^p(\partial\Omega)$ , we have

$$\begin{aligned} S_{2\alpha} S_{3-2\alpha} \phi &= \sum_i \eta_i S_{2\alpha} \kappa_i S_{3-2\alpha} \phi + \sum_i \eta_i S_{2\alpha} (1 - \kappa_i) S_{3-2\alpha} \phi \\ &:= I_1 \phi + I_2 \phi. \end{aligned}$$

Note that for  $P \in \Delta(P_i, 2\epsilon)$ ,  $\kappa_i(\cdot) \Gamma_{2\alpha}(P - \cdot)$  has no singularity in  $\partial\Omega \setminus \Delta(P_i, 3\epsilon)$  and so  $I_2 \phi$  is a smooth function.

For  $I_1\phi$ , we have

$$\begin{aligned}
I_1\phi(P) &= \sum_i \eta_i S_{2\alpha} \kappa_i S_{3-2\alpha} \lambda_i \phi(P) + \sum_i \eta_i S_{2\alpha} \kappa_i S_{3-2\alpha} (1 - \lambda_i) \phi(P) \\
&= \sum_i \eta_i(P) \int_{\partial\Omega} \Gamma_{2\alpha}(P - Z) \kappa_i(Z) \int_{\Delta(P_i, 9\epsilon)} \lambda_i(Q) \Gamma_{3-2\alpha}(Z - Q) \phi(Q) dQ dZ \\
&\quad + \sum_i \eta_i(P) \int_{\partial\Omega} \Gamma_{2\alpha}(P - Z) \kappa_i(Z) \int_{\partial\Omega \setminus \Delta(P_i, 9\epsilon)} (1 - \lambda_i(Q)) \Gamma_{3-2\alpha}(Z - Q) \phi(Q) dQ dZ \\
&:= \sum_i I_{11}^i \phi(P) + \sum_i I_{12}^i \phi(P) \\
&:= I_{11} \phi(P) + I_{12} \phi(P).
\end{aligned}$$

For fixed  $Z \in \Delta(P_i, 7\epsilon)$ ,  $\Gamma_{3-2\alpha}(Z - \cdot)$  has no singularity in  $\partial\Omega \setminus \Delta(P_i, 9\epsilon)$  and so  $\kappa_i(Z) \int_{\partial\Omega \setminus \Delta(P_i, 9\epsilon)} (1 - \lambda_i(Q)) \Gamma_{3-2\alpha}(Z - Q) \phi(Q) dQ$  is a smooth function in  $\partial\Omega$ . Hence, by Proposition 3.1,  $I_{12}\phi$  is a smooth function.

Similarly, we decompose  $S_2\phi$  as

$$S_2\phi(P) = J_{11}\phi(P) + J_{12}\phi(P) + J_2\phi(P),$$

where  $J_{12}\phi$ ,  $J_2\phi$  are smooth functions and

$$J_{11}\phi(P) = \sum_i J_{11}^i \phi(P)$$

with  $J_{11}^i \phi(P) := \eta_i(P) \int_{\Delta(P_i, 9\epsilon)} \Gamma_2(P - Q) \lambda_i(Q) \phi(Q) dQ$ .

For  $I_{11}\phi$ ,  $J_{11}\phi$ , we fix  $i$ . Using the translation and rotation, we assume that  $P_i = 0$  and there is  $\Psi : B'(0, 9\epsilon) \rightarrow \mathbb{R}$  with

$$|\Psi_i(x')| < c|x'|^2 < c\epsilon^2, \quad |\nabla \Psi_i(x')| < c|x'| < c\epsilon, \quad \text{for } x' \in B'(0, 9\epsilon) \quad (4.3)$$

such that for  $Q \in \Delta(P_i, 9\epsilon) := \Delta_{9\epsilon}^i$ ,  $Q$  is represented by  $Q = (y', \Psi(y'))$  for some  $y' \in B'(0, 9\epsilon) := B'(9\epsilon)$ . Let  $P = (x', \Psi(x'))$ ,  $x' \in B'(0, 2\epsilon)$ . Then, we have

$$\begin{aligned}
I_{11}^i \phi(P) &= \eta_i(P) \int_{\Delta_{9\epsilon}^i} \lambda_i(Q) \phi(Q) \int_{\partial\Omega} \kappa_i(Z) \Gamma_{2\alpha}(P - Z) \Gamma_{3-2\alpha}(Q - Z) dZ dQ \\
&= \eta_i(P) \int_{B'(9\epsilon)} \lambda_i(y', \Psi(y')) \phi(y', \Psi(y')) \sqrt{1 + |\nabla \Psi(y')|^2} \times \\
&\quad \int_{B'(7\epsilon)} \kappa_i(z', \Psi(z')) \Gamma_{2\alpha}(x' - z', \Psi(x') - \Psi(z')) \Gamma_{3-2\alpha}(y' - z', \Psi(y') - \Psi(z')) \sqrt{1 + |\nabla \Psi(z')|^2} dz' dy'
\end{aligned}$$

and

$$J_{11}^i \phi(P) := \eta_i(P) \int_{B'(9\epsilon)} \Gamma_2(x' - y', \Psi(x') - \Psi(y')) \lambda_i(y', \Psi(y')) \phi(y', \Psi(y')) \sqrt{1 + |\nabla \Psi(y')|^2} dy'.$$

Let

$$\begin{aligned} I_{111}^i \phi(P) &:= \eta_i(P) \int_{B'(9\epsilon)} \lambda_i(y', 0) \phi(y', \Psi(y')) \int_{B'(7\epsilon)} \kappa_i(z', 0) \Gamma_{2\alpha}(x' - z', 0) \Gamma_{3-2\alpha}(y' - z', 0) dz' dy', \\ J_{111}^i \phi(P) &:= \eta_i(P) \int_{B'(9\epsilon)} \lambda_i(y', 0) \Gamma_2(x' - y', 0) \phi(y', \Phi(y')) dy'. \end{aligned}$$

Note that by (4.3), we have

$$\|I_{11}^i - I_{111}^i\|_{L^p(\Delta_{9\epsilon}^i) \rightarrow H_p^1(\Delta_{2\epsilon}^i)}, \quad \|J_{11}^i - J_{111}^i\|_{L^p(\Delta_{9\epsilon}^i) \rightarrow H_p^1(\Delta_{2\epsilon}^i)} \leq c\epsilon.$$

(See [6]).

It is well-known

$$\int_{\mathbb{R}^{n-1}} \Gamma_{2\alpha}(x' - z', 0) \Gamma_{3-2\alpha}(y' - z', 0) dz' = \Gamma_2(x' - y', 0).$$

(See Section 5.1 in [17].) Hence, we have

$$\begin{aligned} \Gamma_2(x' - y', 0) &= \int_{\mathbb{R}^{n-1}} \kappa_i(z', 0) \Gamma_{2\alpha}(x' - z', 0) \Gamma_{3-2\alpha}(y' - z', 0) dz' \\ &\quad + \int_{\mathbb{R}^{n-1}} (1 - \kappa_i(z', 0)) \Gamma_{2\alpha}(x' - z', 0) \Gamma_{3-2\alpha}(y' - z', 0) dz' \\ &= \int_{\mathbb{R}^{n-1}} \kappa_i(z', 0) \Gamma_{2\alpha}(x' - z', 0) \Gamma_{3-2\alpha}(y' - z', 0) dz' + k_i(x', y'). \end{aligned}$$

Hence, we have that

$$I_{111}^i \phi(P) - J^i(111) \phi(P) = \eta_i(P) \int_{B'(9\epsilon)} \lambda_i(y', 0) \phi(y', \Psi(y')) k_i(x', y') dy'$$

is a smooth function. Let  $T^1 = \sum_i (I_{11}^i - I_{111}^i) + \sum_i (J_{11}^i - J_{111}^i)$  and  $T_2 = I_2 + J_2 + I_{12} + J_{12} + \sum_i (I_{111}^i - J^i(111))$ . Then,  $S_{2\alpha} S_{3-2\alpha} = S_2 + T^1 + T^2$  such that  $T^2$  is a smooth function and

$$\begin{aligned} \|T^1 \phi\|_{H_p^1(\partial\Omega)} &\leq c \sum_i (\|(I_{11}^i - I_{111}^i) \phi\|_{H_p^1(\partial\Omega)} + \|(J_{11}^i - J_{111}^i) \phi\|_{H_p^1(\partial\Omega)}) \\ &\leq c\epsilon \sum_i \|\phi\|_{L^p(\Delta_{9\epsilon}^i)} \leq c\epsilon \|\phi\|_{L^p(\partial\Omega)}. \end{aligned}$$

Hence, we complete the proof of Lemma 4.1.  $\square$

**Proof Theorem 1.1.**

(1). In the case of  $p \geq p_0 := \frac{2(n-1)}{n-2+2\alpha}$  ( $p_0 < 2$ ). To show the injectivity, suppose that  $S_{2\alpha} \phi = 0$  for  $\phi \in L^p(\partial\Omega)$ . By the Hölder inequality and Sobolev imbedding, we have  $L^p(\partial\Omega) \subset L^{p_0}(\partial\Omega) \subset H_2^{-\alpha+\frac{1}{2}}(\partial\Omega)$ . Hence, by the bijectivity of  $S_{2\alpha} : H_2^{-\alpha+\frac{1}{2}}(\partial\Omega) \rightarrow H_2^{\alpha-\frac{1}{2}}(\partial\Omega)$  (see [5]), we have  $\phi = 0$ . Hence,  $S_{2\alpha} : L^p(\partial\Omega) \rightarrow H_p^{2\alpha-1}(\partial\Omega)$  is bijective for  $p \geq p_0$ .

To show that  $S_{2\alpha} : L^p(\partial\Omega) \rightarrow H_p^{2\alpha-1}(\partial\Omega)$  is surjective, let  $f \in H_p^{2\alpha-1}(\partial\Omega)$ . By the Sobolev imbedding and the Hölder inequality, we have  $H_p^{2\alpha-1}(\partial\Omega) \subset H_{p_0}^{2\alpha-1}(\partial\Omega) \subset H_2^{\alpha-\frac{1}{2}}(\partial\Omega)$ . By the bijectivity of  $S_{2\alpha} : H_2^{-\alpha+\frac{1}{2}}(\partial\Omega) \rightarrow H_2^{\alpha-\frac{1}{2}}(\partial\Omega)$ , there is  $\phi \in H_2^{-\alpha+\frac{1}{2}}(\partial\Omega)$  such that  $S_{2\alpha} \phi = f$ . Note that by the Lemma 4.1, we get that  $S_{3-2\alpha} S_{2\alpha} = S_2 + T^1 + T^2$ , where  $\|T^1\|_{L^p(\partial\Omega) \rightarrow H_p^1(\partial\Omega)} \leq \epsilon$  and  $T^2 : H_2^{-1}(\partial\Omega) \rightarrow H_2^1(\partial\Omega)$  is bounded. Then, by the Lemma 4.1, we obtain that  $(S_2 + T^1) \phi =$

$S_{3-2\alpha}S_{2\alpha}\phi - T^2\phi \in H_p^1(\partial\Omega)$ . Taking  $\epsilon > 0$  sufficiently small such that  $S_2 + T^1 : L^p(\partial\Omega) \rightarrow H_p^1(\partial\Omega)$  is bijective (see (1) of the Remark 4.2), we obtain that  $\phi \in L^p(\partial\Omega)$ . This implies that  $S_{2\alpha} : L^p(\partial\Omega) \rightarrow H_p^{2\alpha-1}(\partial\Omega)$  is surjective. Hence, we complete the proof of the bijectivity of  $S_{2\alpha} : L^p(\partial\Omega) \rightarrow H_p^{2\alpha-1}(\partial\Omega)$  for  $p \geq \frac{2(n-1)}{n-2+2\alpha}$ .  $\square$

**Remark 4.3.** (1) Note that the dual operator  $S_{2\alpha}^* : H_{p'}^{-2\alpha+1}(\partial\Omega) \rightarrow L^{p'}(\partial\Omega)$  of  $S_{2\alpha} : L^p(\partial\Omega) \rightarrow H_p^{2\alpha-1}(\partial\Omega)$  are same with the operator  $S_{2\alpha} : H_{p'}^{-2\alpha+1}(\partial\Omega) \rightarrow L^{p'}(\partial\Omega)$ , where  $\frac{1}{p} + \frac{1}{p'} = 1$  by the section 3. Hence, by the property of the dual operator, we get that  $S_{2\alpha} : H_p^{-2\alpha+1}(\partial\Omega) \rightarrow L^p(\partial\Omega)$  is bijective for  $1 < p \leq \frac{2(n-1)}{n-2\alpha} = p_0'$ .

(2) In the Lemma 4.1,  $S_{3-2\alpha}S_{2\alpha}$  is sum of a bijective operator  $S_2 + T^1$  and a compact operator  $T^2$  and so  $S_{3-2\alpha}S_{2\alpha}$  is the Fredholm operator with index zero. Since  $S_{2\alpha} : L^p(\partial\Omega) \rightarrow H_p^{2\alpha-1}(\partial\Omega)$ ,  $S_{3-2\alpha} : H_p^{2\alpha-1}(\partial\Omega) \rightarrow H_p^1(\partial\Omega)$  are injective,  $S_{3-2\alpha}S_{2\alpha}$  is injective and so by the Fredholm operator theorem,  $S_{3-2\alpha}S_{2\alpha} : L^p(\partial\Omega) \rightarrow H_p^1(\partial\Omega)$  is bijective. This implies that  $S_{3-2\alpha} : H_p^{2\alpha-1}(\partial\Omega) \rightarrow H_p^1(\partial\Omega)$  is bijective for  $p \geq \frac{2(n-1)}{(n-2+2\alpha)}$ .

(2). In the case of  $1 < p < 2$ . Now, we will show that  $S_{2\alpha} : H_q^{-2\alpha+1}(\partial\Omega) \rightarrow L^q(\partial\Omega)$  is surjective for  $q = \frac{p}{p-1} > 2$ . Let  $f \in L^q(\partial\Omega)$ . By the Hölder inequality,  $L^q(\partial\Omega) \subset L^2(\partial\Omega)$  and by the bijectivity of  $S_{2\alpha} : H_2^{-2\alpha+1}(\partial\Omega) \rightarrow L^2(\partial\Omega)$  (see (1) of Remark 4.3), there is  $\phi \in H_2^{-2\alpha+1}(\partial\Omega)$  such that  $S_{2\alpha}\phi = f$ . Then,

$$S_{3-2\alpha}S_{2\alpha}\phi = S_{3-2\alpha}f \in H_q^{2-2\alpha}(\partial\Omega).$$

By (3) of Remark 4.2, we obtain that  $(S_2 + T^1)\phi \in H_q^{2-2\alpha}(\partial\Omega)$ . Since  $S_2 + T^1 : H_q^{-2\alpha+1}(\partial\Omega) \rightarrow H_q^{2-2\alpha}(\partial\Omega)$  is bijective (see (3) of Remark 4.2), we obtain that  $\phi \in H_q^{-2\alpha+1}(\partial\Omega)$ . This implies that  $S_{2\alpha} : H_q^{-2\alpha+1}(\partial\Omega) \rightarrow L^q(\partial\Omega)$  is surjective.

By the property of the dual operator, we have that  $S_{2\alpha}^* : L^p(\partial\Omega) \rightarrow H_p^{2\alpha-1}(\partial\Omega)$ ,  $1 < p < 2$  are injective. Since  $S_{2\alpha}^* = S_{2\alpha}$ ,  $S_{2\alpha} : L^p(\partial\Omega) \rightarrow H_p^{2\alpha-1}(\partial\Omega)$  is injective and so  $S_{2\alpha} : H_p^{2\alpha-1}(\partial\Omega) \rightarrow H_p^1(\partial\Omega)$  is injective. Hence, we get  $S_{2\alpha}S_{3-2\alpha} : L^p(\partial\Omega) \rightarrow H_p^1(\partial\Omega)$ ,  $1 < p < 2$  are injective.

Note that in (3) of Remark 4.2,  $S_{2\alpha}S_{3-2\alpha}$  is the sum of a bijective operator and a compact operator. Hence, by the Fredholm theorem,  $S_{2\alpha}S_{3-2\alpha} : L^p(\partial\Omega) \rightarrow H_p^1(\partial\Omega)$  is bijective.

To show that  $S_{2\alpha} : L^p(\partial\Omega) \rightarrow H_p^{2\alpha-1}(\partial\Omega)$  is surjective for  $1 < p < 2$ , let us  $f \in H_p^{2\alpha-1}(\partial\Omega)$ . Then, we have  $S_{3-2\alpha}f \in H_p^1(\partial\Omega)$ . Since,  $S_{3-2\alpha}S_{2\alpha} : L^p(\partial\Omega) \rightarrow H_p^1(\partial\Omega)$  is bijective, there is  $\phi \in L^p(\partial\Omega)$  such that  $S_{3-2\alpha}S_{2\alpha}\phi = S_{3-2\alpha}f$ . Since  $S_{3-2\alpha}$  is injective, we get  $S_{2\alpha}\phi = f$  and so  $S_{2\alpha} : L^p(\partial\Omega) \rightarrow H_p^{2\alpha-1}(\partial\Omega)$  is bijective.  $\square$

**Corollary 4.4.** *Let  $\frac{1}{2} < \alpha < 1$  and  $1 < p < \infty$ . For  $-2\alpha + 1 \leq s \leq 2 - 2\alpha$ ,*

$$\begin{aligned} S_{2\alpha} : H_p^s(\partial\Omega) &\rightarrow H_p^{s+2\alpha-1}(\partial\Omega), \\ S_{2\alpha} : B_p^s(\partial\Omega) &\rightarrow B_p^{s+2\alpha-1}(\partial\Omega) \end{aligned}$$

*are bijective.*

*Proof.* In the proof of the Theorem 1.1, we have  $S_{3-2\alpha} : L^p(\partial\Omega) \rightarrow H_p^{2-2\alpha}(\partial\Omega)$ ,  $S_{2\alpha} : H_p^{2-2\alpha}(\partial\Omega) \rightarrow H_p^1(\partial\Omega)$  are injective and so  $S_{2\alpha}S_{3-2\alpha} : L^p(\partial\Omega) \rightarrow H_p^1(\partial\Omega)$  is injective. Since  $S_{2\alpha}S_{3-2\alpha} : L^p(\partial\Omega) \rightarrow H_p^1(\partial\Omega)$  is Fredholm operator with index zero, we get  $S_{2\alpha}S_{3-2\alpha} : L^p(\partial\Omega) \rightarrow H_p^1(\partial\Omega)$  is bijective. This implies

$$S_{2\alpha} : H_p^{2-2\alpha}(\partial\Omega) \rightarrow H_p^1(\partial\Omega) \quad \text{is bijective.} \quad (4.4)$$

By the dual operator property and the fact of  $S_{2\alpha}^* = S_{2\alpha}$ , we have

$$S_{2\alpha} : H_p^{-1}(\partial\Omega) \rightarrow H_p^{-2+2\alpha}(\partial\Omega) \quad \text{is bijective.} \quad (4.5)$$

Using (4.4), (4.5) and the properties of the real interpolation and complex interpolation, we obtain the corollary. □

## 5. PROOF OF THEOREM 1.2

We introduce a Riesz potential  $I_{2\alpha}$ ,  $0 < 2\alpha < n$ , by

$$I_{2\alpha}f(x) = c(n, \alpha) \int_{\mathbb{R}^n} \frac{1}{|x-y|^{n-2\alpha}} f(y) dy \quad \text{for } \psi \in C_c^\infty(\mathbb{R}^n),$$

where  $c(n, \alpha) := \frac{(2\pi)^{2\alpha} \Gamma(\frac{n}{2} - \alpha)}{\pi^{\frac{n}{2}} 2^{2\alpha} \Gamma(\alpha)}$ .

The following proposition is well known fact and will be useful in the subsequent estimates (see chapter 5 of [17]).

**Proposition 5.1.** 1). *Let  $1 < p < q < \infty$ ,  $\frac{1}{q} = \frac{1}{p} - \frac{2\alpha}{n}$ . Then*

$$I_{2\alpha} : L^p(\mathbb{R}^n) \rightarrow L^q(\mathbb{R}^n)$$

*is bounded.*

2). *Let  $1 < p < \infty$  and  $s \in \mathbb{R}$ . Then, the following operators are bounded.*

$$I_{2\alpha} : H_p^s(\mathbb{R}^n) \rightarrow H_p^{s+2\alpha}(\mathbb{R}^n), \quad I_{2\alpha} : B_p^s(\mathbb{R}^n) \rightarrow B_p^{s+2\alpha}(\mathbb{R}^n).$$

**Remark 5.2.** *Let  $B_R$  be the open ball in  $\mathbb{R}^n$  centered at the origin with radius  $R$ . Then, by proposition 5.1, the following operator is bounded.*

$$\tilde{I}_{2\alpha} : B_{p0}^s(B_R) \rightarrow B_p^{s+2\alpha}(B_R), \quad \tilde{I}_{2\alpha}f(x) = \int_{\mathbb{R}^n} \Gamma_{2\alpha}(x-y)f(y)dy, \quad f \in B_{p0}^s(B_R), \quad s \in \mathbb{R}.$$

**Proof of (1.4)** Let  $-2\alpha + 1 - \frac{1}{p} < s < 0$ . Let  $\phi \in C^1(\partial\Omega)$  and  $f \in C_c^\infty(B_R)$ . Then, we have

$$\int_{\mathbb{R}^n} f(x) \mathcal{S}_{2\alpha} \phi(x) dx = \int_{\partial\Omega} \phi(P) \tilde{I}_{2\alpha} f(P) dP.$$

Since  $C^1(\partial\Omega)$  is dense subspace of  $B_p^s(\partial\Omega)$  and  $C_c^\infty(B_R)$  is dense subspace of  $B_{q0}^{-s-2\alpha+\frac{1}{q}}(B_R)$ , we have

$$\langle f, \mathcal{S}_{2\alpha} \phi \rangle_{(B_{q0}^{-s-2\alpha+\frac{1}{q}}(B_R), B_p^{s+2\alpha-1+\frac{1}{p}}(B_R))} = \langle \phi, \tilde{I}_{2\alpha} f \rangle_{(B_p^s(\partial\Omega), B_q^{-s}(\partial\Omega))},$$

where  $\frac{1}{p} + \frac{1}{q} = 1$ . Then, we have

$$\begin{aligned} \langle f, \mathcal{S}_{2\alpha} \phi \rangle_{(B_{q0}^{-s-2\alpha+\frac{1}{q}}(B_R), B_p^{s+2\alpha-1+\frac{1}{p}}(B_R))} &\leq \|\phi\|_{B_p^s(\partial\Omega)} \|\tilde{I}_{2\alpha} f\|_{B_q^{-s}(\partial\Omega)} \\ &\leq c \|\phi\|_{B_p^s(\partial\Omega)} \|\tilde{I}_{2\alpha} f\|_{B_q^{-s+\frac{1}{q}}(B_R)} \\ &\leq c \|\phi\|_{B_p^s(\partial\Omega)} \|f\|_{B_{q0}^{-s-2\alpha+\frac{1}{q}}(B_R)}. \end{aligned}$$

Hence, we have

$$\begin{aligned} \|\mathcal{S}_{2\alpha} \phi\|_{B_p^{s+2\alpha-1+\frac{1}{p}}(B_R)} &= \sup_{\|f\|_{B_{q0}^{-s-2\alpha+\frac{1}{q}}(B_R)}=1} |\langle f, \mathcal{S}_{2\alpha} \phi \rangle_{(B_{q0}^{-s-2\alpha+\frac{1}{q}}(B_R), B_p^{s+2\alpha-1+\frac{1}{p}}(B_R))}| \\ &\leq c \|\phi\|_{B_p^s(\partial\Omega)}. \end{aligned}$$

We complete the proof of (1.4).  $\square$

**Proof of (1.5).** For  $\phi \in B_p^s(\partial\Omega)$ ,  $-2\alpha + 1 - \frac{1}{p} < s < 0$ , let us  $u$  be a the layer potential of  $\phi$  defined by (1.3). Note that  $u$  is in  $C^\infty(\mathbb{R}^n \setminus \partial\Omega)$  and for large  $|x|$ , we have

$$|D^\beta u(x)| \leq \|\phi\|_{B_p^s(\partial\Omega)} \|D^\beta \Gamma_{2\alpha}(x - \cdot)\|_{B_{p'}^{-s}(\partial\Omega)} \leq c \|\phi\|_{B_p^s(\partial\Omega)} \frac{1}{|x|^{n-2\alpha+|\beta|}}, \quad (5.1)$$

where  $\frac{1}{p} + \frac{1}{p'} = 1$ .

Let  $B_R$  be an open ball whose center is origin and radius is  $R \geq 2$  such that  $\Omega \subset B_{\frac{1}{3}R}$ . We divide the left-hand side of (1.5) into three parts

$$\begin{aligned} A_1 &= \int_{|x| \leq R} \int_{|y| \leq R} \frac{|D_x^k u(x) - D_y^k u(y)|^p}{|x-y|^{n+p(s+2\alpha-k-1+\frac{1}{p})}} dy dx, \\ A_2 &= 2 \int_{|x| \leq R} \int_{|y| \geq R} \frac{|D_x^k u(x) - D_y^k u(y)|^p}{|x-y|^{n+p(s+2\alpha-k-1+\frac{1}{p})}} dy dx, \\ A_3 &= \int_{|x| \geq R} \int_{|y| \geq R} \frac{|D_x^k u(x) - D_y^k u(y)|^p}{|x-y|^{n+p(s+2\alpha-k-1+\frac{1}{p})}} dy dx. \end{aligned} \quad (5.2)$$

By (1.4),  $A_1$  is dominated by  $\|\phi\|_{B_p^s(\partial\Omega)}^p$ . For  $|x| \leq R$  and  $|y| \geq 2R$ , we get that  $|x-y| \geq |y| - |x| \geq |y| - R \geq \frac{1}{2}|y|$ . Note that by (5.1), for  $|y| \geq 2R$ , we have that  $|D_y^k u(y)| \leq c|y|^{-n+2\alpha-k} \|\phi\|_{B_p^s(\partial\Omega)}^2$ .

Hence, by (1.4), we have

$$\begin{aligned}
A_2 &\leq 2 \int_{|x| \leq R} \int_{R \leq |y| \leq 2R} \frac{|D_x^k u(x) - D_y^k u(y)|^p}{|x - y|^{n+p(s+2\alpha-k-1+\frac{1}{p})}} dy dx \\
&\quad + 2^{n+2s+2} \int_{|x| \leq R} \int_{|y| \geq 2R} \frac{|D_x^k u(x)|^p + |D_y^k u(y)|^p}{|y|^{n+p(s+2\alpha-k-1+\frac{1}{p})}} dy dx \\
&\leq c_R \|u\|_{B_p^{s+2\alpha-1+\frac{1}{p}}(B(2R))}^p + c \|\phi\|_{B_p^s(\partial\Omega)}^p \int_{|x| \leq R} \int_{|y| \geq 2R} \frac{1}{|y|^{n+p(s-1+\frac{1}{p}+n)}} dy dx \\
&\leq c_R \|\phi\|_{B_p^s(\partial\Omega)}^p.
\end{aligned}$$

We divide  $A_3$  into two parts;

$$\begin{aligned}
&\int_{|x| \geq R} \int_{|y| \geq R, |x-y| \leq \frac{1}{2}|x|} \frac{|D_x^k u(x) - D_y^k u(y)|^p}{|x-y|^{n+p(s+2\alpha-k-1+\frac{1}{p})}} dy dx \\
&\quad + \int_{|x| \geq R} \int_{|y| \geq R, |x-y| \geq \frac{1}{2}|x|} \frac{|D_x^k u(x) - D_y^k u(y)|^p}{|x-y|^{n+p(s+2\alpha-k-1+\frac{1}{p})}} dy dx.
\end{aligned} \tag{5.3}$$

Applying the mean-value theorem, for  $|x| \geq R$ ,  $|x-y| \leq \frac{1}{2}|x|$ , there is a  $\xi$  between  $x$  and  $y$  such that  $D_x^k u(x) - D_y^k u(y) = D^{k+1}u(\xi) \cdot (x-y)$ . Note that  $|x-\xi| \leq \frac{1}{2}|x|$  and hence  $|\xi| \geq \frac{1}{2}|x| \geq \frac{1}{2}R$ . Since  $s+2\alpha-k-2+\frac{1}{p} < 0$  and  $p > \frac{n-1}{n+s-1}$ , by (5.1), the first term of (5.3) is dominated by

$$\begin{aligned}
&\int_{|x| \geq R} \int_{|y| \geq R, |x-y| \leq \frac{1}{2}|x|} \frac{|D^{k+1}u(\xi)|^p}{|x-y|^{n+p(s+2\alpha-k-1+\frac{1}{p})-p}} dy dx \\
&\leq c \|\phi\|_{B_p^s(\partial\Omega)}^p \int_{|x| \geq R} \frac{1}{|x|^{pn-2p\alpha+(k+1)p}} \int_{|x-y| \leq \frac{1}{2}|x|} \frac{1}{|x-y|^{n+p(s+2\alpha-k-2+\frac{1}{p})}} dy dx \\
&\leq c \|\phi\|_{B_p^s(\partial\Omega)}^p \int_{|x| \geq R} \frac{1}{|x|^{p(n+s-1)+1}} dx \\
&= cR^{-p(n+s-1)-1+n} \|\phi\|_{B_p^s(\partial\Omega)}^p.
\end{aligned}$$

Since  $|x|, |y| \geq R$ , by (5.1), the second term of (5.3) is dominated by

$$\begin{aligned}
&\int_{|x| \geq R} \int_{|y| \geq R, |x-y| \geq \frac{1}{2}|x|} \frac{|D_x^k u(x)|^p + |D_y^k u(y)|^p}{|x-y|^{n+p(s+2\alpha-k-1+\frac{1}{p})}} dy dx \\
&\leq \|\phi\|_{B_p^s(\partial\Omega)}^p \int_{|x| \geq R} \frac{1}{|x|^{p(n-2\alpha+k)}} \int_{|y| \geq R, |x-y| \geq \frac{1}{2}|x|} \frac{1}{|x-y|^{n+p(s+2\alpha-k-1+\frac{1}{p})}} dy dx \\
&\quad + \|\phi\|_{B_p^s(\partial\Omega)}^p \int_{|x| \geq R} \int_{|y| \geq R, |x-y| \leq \frac{1}{2}|x|} \frac{1}{|x-y|^{n+p(s+2\alpha-k-1+\frac{1}{p})}} \frac{1}{|y|^{p(n-2\alpha+k)}} dy dx.
\end{aligned} \tag{5.4}$$

Since  $p > \frac{n-1}{n+s-1}$ , the second term of right-hand side of (5.4) is dominated by  $R^{-p(n+s-1)-1+n} \|\phi\|_{B_p^s(\partial\Omega)}^p$ .

Note that

$$\begin{aligned}
&\int_{|x| \geq R} \int_{R \leq |y| \leq 2|x|} \frac{1}{|x|^{n+p(s+2\alpha-k-1+\frac{1}{p})}} \frac{1}{|y|^{pn-2p\alpha+kp}} dy dx \\
&\leq c \begin{cases} R^{-pn+2p\alpha+n} \int_{|x| \geq R} \frac{1}{|x|^{n+p(s+2\alpha-k-1+\frac{1}{p})}} dx, & pn-2p\alpha+kp > n, \\ \int_{|x| \geq R} \frac{\ln|x|}{|x|^{n+p(s+2\alpha-k-1+\frac{1}{p})}} dx, & pn-2p\alpha+kp = n, \\ \int_{|x| \geq R} \frac{1}{|x|^{p(n+s-k-1)+1}} dx, & pn-2p\alpha+kp < n \end{cases} \\
&\leq cR^{-p(n-k-1+s)-1+n} \ln R.
\end{aligned}$$

Then, since  $p > \frac{n-1}{n+s-1}$ , the first term of right-hand side of (5.4) is dominated by

$$\begin{aligned}
& \|\phi\|_{B_p^s(\partial\Omega)}^p \int_{|x| \geq R} \int_{|y| \geq R, |x-y| \geq \frac{1}{2}|x|} \frac{1}{|x-y|^{n+p(s+2\alpha-k-1+\frac{1}{p})}} \frac{1}{|y|^{pn-2p\alpha+kp}} dy dx \\
& \leq c \|\phi\|_{B_p^s(\partial\Omega)}^p \left( \int_{|x| \geq R} \int_{R \leq |y| \leq 2|x|} \frac{1}{|x|^{n+p(s+2\alpha-k-1+\frac{1}{p})}} \frac{1}{|y|^{pn-2p\alpha+kp}} dy dx \right. \\
& \quad \left. + \int_{|x| \geq R} \int_{|y| \geq 2|x|} \frac{1}{|y|^{n+p(n+s-1)+1}} dy dx \right) \\
& \leq c R^{-p(n-1+s)-1+n} \ln R \|\phi\|_{B_p^s(\partial\Omega)}^p.
\end{aligned}$$

Therefore, we showed that  $A_1 + A_2 + A_3 \leq c_R \|\phi\|_{B_p^s(\partial\Omega)}^p$  and hence showed (1.5).  $\square$

## 6. PROOF OF THEOREM 1.4

**Theorem 6.1.** *Let  $1 - 2\alpha - \frac{1}{p} < s < 0$ . For  $\phi \in B_p^s(\partial\Omega)$ , let us  $u = \mathcal{S}_{2\alpha}\phi$  be a layer potential defined in (1.3). Then, we have*

(1)

$$\hat{u}(\xi) = |\xi|^{-2\alpha} < \phi, e^{2\pi i \xi \cdot} >. \quad (6.1)$$

(2)  $u$  is a weak solution of

$$\Delta^\alpha u = 0, \quad \text{in } \mathbb{R}^n \setminus \partial\Omega. \quad (6.2)$$

*Proof.* For the proof of (6.1), let us  $\phi \in C^2(\partial\Omega)$  and  $\psi \in C_c^\infty(\mathbb{R}^n)$ . Then, we have

$$\begin{aligned}
\int_{\mathbb{R}^n} u(x) \psi(x) dx &= c(n, s) \int_{\partial\Omega} \phi(Q) \int_{\mathbb{R}^n} \frac{1}{|x-Q|^{n-2\alpha}} \psi(x) dx dQ \\
&= \int_{\partial\Omega} \phi(Q) \int_{\mathbb{R}^n} |\xi|^{-2\alpha} e^{2\pi i \xi \cdot Q} \overline{\hat{\psi}(\xi)} d\xi dQ \\
&= \int_{\mathbb{R}^n} \overline{\hat{\psi}(\xi)} |\xi|^{-2\alpha} \int_{\partial\Omega} \phi(Q) e^{2\pi i \xi \cdot Q} dQ d\xi.
\end{aligned}$$

Hence, we get

$$\hat{u}(\xi) = |\xi|^{-2\alpha} \int_{\partial\Omega} \phi(Q) e^{2\pi i \xi \cdot Q} dQ.$$

Since  $C^2(\partial\Omega)$  is dense in  $B_p^s(\partial\Omega)$ , we get (6.1) for all  $\phi \in B_p^s(\partial\Omega)$ .

For the proof (6.2), suppose that  $\phi \in C^2(\partial\Omega)$  and  $\psi \in C^\infty(\mathbb{R}^n \setminus \partial\Omega)$ , then, by (6.1),

$$\begin{aligned}
\int_{\mathbb{R}^n} u(x) \Delta^\alpha \psi(x) dx &= \int_{\mathbb{R}^n} |\xi|^{2\alpha} \hat{u}(\xi) \overline{\hat{\psi}(\xi)} d\xi \\
&= \int_{\mathbb{R}^n} \overline{\hat{\psi}(\xi)} \int_{\partial\Omega} e^{-2\pi i \xi \cdot Q} \phi(Q) dQ d\xi \\
&= \int_{\partial\Omega} \phi(Q) \overline{\int_{\mathbb{R}^n} e^{2\pi i \xi \cdot Q} \hat{\psi}(\xi) d\xi} dQ \\
&= \int_{\partial\Omega} \phi(Q) \psi(Q) dQ \\
&= 0
\end{aligned} \quad (6.3)$$

Since  $\Delta^t : \dot{B}_p^s(\mathbb{R}^n) \rightarrow \dot{B}_p^{s-2t}(\mathbb{R}^n)$  is isomorphism, we have

$$\begin{aligned} \left| \int_{\mathbb{R}^n} u(x) \Delta^\alpha \psi(x) dx \right| &= \left| \int_{\mathbb{R}^n} \Delta^{\frac{1}{2}(s+2\alpha-1+\frac{1}{p})} u(x) \Delta^{\frac{1}{2}(-s+1-\frac{1}{p})} \psi(x) dx \right| \\ &\leq \left\| \Delta^{\frac{1}{2}(s+2\alpha-1+\frac{1}{p})} u \right\|_{\dot{B}_p^0(\mathbb{R}^n)} \left\| \Delta^{\frac{1}{2}(-s+1-\frac{1}{p})} \psi \right\|_{\dot{B}_{p'}^0(\mathbb{R}^n)} \\ &\leq \|u\|_{\dot{B}_p^{s+2\alpha-1+\frac{1}{p}}(\mathbb{R}^n)} \|\psi\|_{\dot{B}_{p'}^{-s+1-\frac{1}{p}}(\mathbb{R}^n)} \\ &\leq c \|\phi\|_{B_p^s(\partial\Omega)} \|\psi\|_{B_{p'}^{-s+1-\frac{1}{p}}(\mathbb{R}^n)}. \end{aligned}$$

Let  $\phi_k \in C^2(\partial\Omega)$  such that  $\phi_k \rightarrow \phi$  in  $B_p^s(\partial\Omega)$  and  $u_k = \mathcal{S}\phi_k$ . Then, we get

$$\left| \int_{\mathbb{R}^n} (u_k(x) - u(x)) \Delta^\alpha \psi(x) dx \right| \leq c \|\phi_k - \phi\|_{B_p^s(\partial\Omega)} \|\psi\|_{B_{p'}^{-s+1-\frac{1}{p}}(\mathbb{R}^n)} \rightarrow 0, \quad k \rightarrow \infty.$$

Hence, since  $C^2(\partial\Omega)$  is the dense subspace of  $B_p^s(\partial\Omega)$ , (6.3) holds for  $\phi \in B_p^s(\partial\Omega)$  and so we get (6.2) for all  $\phi \in B_p^s(\partial\Omega)$ .  $\square$

**Proof of Theorem 1.4.** By the Corollary 4.4, we get that  $S_{2\alpha} : B_p^{t-2\alpha+1}(\partial\Omega) \rightarrow B_p^t(\partial\Omega)$ ,  $0 < t < 1$ ,  $1 < p < \infty$  are bijective.

To show the existence of solution, let  $g \in B_p^t(\partial\Omega)$ . By the bijectivity of  $S_{2\alpha} : B_p^{t-2\alpha+1}(\partial\Omega) \rightarrow B_p^t(\partial\Omega)$ , there is a  $\phi \in B_p^{t-2\alpha+1}(\partial\Omega)$  such that  $S_{2\alpha}\phi = g$ . Let  $u = \mathcal{S}_{2\alpha}\phi$  defined by (1.3). Then, by (2) of the Theorem 6.1,  $u$  is a weak solution of (6.2) and by the Theorem 1.2,  $u$  satisfies the equation (1.7). Hence, we complete the proof of the Theorem 1.4.  $\square$

## REFERENCES

- [1] J. Bergh and J. Lofström, *Interpolation spaces, An introduction*, Springer-Verlag, Berlin(1976).
- [2] K. Bogdan and T. Byczkowski, *Potential theory for the  $\alpha$ -stable Schrodinger operator on bounded Lipschitz domains*, Studia Math. **133**, no. 1, 53-92(1999).
- [3] Russell M. Brown, *The method of layer potentials for the heat equation in Lipschitz cylinders*, Amer. J. Math, **111**, no. 2, 339-379(1989).
- [4] Russell M. Brown, *The initial-Neumann problem for the heat equation in Lipschitz cylinders*, Trans. Amer. Math. Soc, **320**, no. 1, 1-52(1990).
- [5] T.K. Chang, *Boundary integral operator for the fractional Laplace equation in a bounded Lipschitz domain*, Integral Equations Operator Theory, **72**, no. 3, 345-361(2012).
- [6] R. R. Coifman, A. McIntosh and Y. Meyer, *L'integrale de Cauchy definit un operateur Borne sur  $L^2$  pour les Courbes Lipschitiennes*, Ann. of Math., **116**, pp. 361-388(1982).
- [7] B. E. J. Dahlberg, C. E. Kenig and G. C. Verchota, *Boundary value problems for the systems of elastostatics in Lipschitz domains*, Duke Math. J, **57**, no. 3, 109-135(1988).
- [8] M. Frazier and B. Jawerth, *Decomposition of Besov spaces*, Indiana Univ. Math. J., **34**, no. 4, 777-799(1985).
- [9] E. Fabes, M. Jodeit and N. Riviere, *Potential techniques for boundary value problems on  $C^1$ -domains*. Acta Math, **141**, no. 3-4, 165-186(1978).
- [10] E. B. Fabes, C. E. Kenig and G. C. Verchota, *The Dirichlet problem for the Stokes system on Lipschitz domains*, Duke Math. J, **57**, no. 3, 769-793(1988).
- [11] E. Fabes, O. Mendez and M. Mitrea, *Boundary layers on Sobolev-Besov spaces and Poisson's equation for the Laplacian in Lipschitz domains*, J. Funct. Anal, **159**, , no. 2, 323-368(1998).
- [12] S. Hofmann and J.L. Lewis,  *$L^2$  solvability and representation by caloric layer potentials in time-varying domains*, Ann. of Math. (2), **144**, no. 2, 349-420(1996).
- [13] D. Jerison and C. E. Kenig, *The inhomogeneous Dirichlet problem in Lipschitz domains*, J. Funct. Anal, **130**, no. 1, 161-219(1995).
- [14] A. Jonsson and H. Wallin, *A Whitney extension theorem in  $L_p$  and Besov spaces*, Ann. Inst. Fourier. Grenoble, **28**, no 1, 139-192(1978).

- [15] M. Mitrea, *The method of layer potentials in electromagnetic scattering theory on nonsmooth domains*, Duke Math. J., **77**, no. 1, 111-133(1995).
- [16] Z. W. Shen, *Boundary value problems for parabolic Lamé systems and a nonstationary linearized system of Navier-Stokes equations in Lipschitz cylinders*, Amer. J. Math, **113**, no. 2(1991), 293-373.
- [17] E. M. Stein, *Singular Integrals and Differentiability Properties of Functions*, Princeton University Press, Princeton, N.J, 1970.
- [18] G. Verchota, *Layer potentials and regularity for the Dirichlet problem for Laplace's equation in Lipschitz domains*, J. of Funct. Anal, **59**(1984), 572-611.
- [19] M. Zähle, *Harmonic calculus on fractals- A measure geometric approach II*, Trans. Amer. Math. Soc, **357**, no. 9(2005), 3407-3423.
- [20] M. Zähle, *Riesz potentials and Liouville operators on fractals*, Potential Anal., **21**, no. 2(2004), 193-208.

DEPARTMENT OF MATHEMATICS, YONSEI UNIVERSITY, SEOUL, 136-701, SOUTH KOREA  
*E-mail address:* `chang7357@yonsei.ac.kr`